

Sexual health service needs by gender and sexual orientation among clients in British Columbia during the first few months of the COVID-19 pandemic response

Aidan Ablona¹, <u>Hsiu-Ju Chang</u>¹, Travis Salway², Daniel Grace³, Catherine Worthington⁴, Mark Gilbert^{1,5}

¹BC Centre for Disease Control, ²Simon Fraser University, ³University of Toronto, ⁴University of Victoria, ⁵University of British Columbia

CAHR Virtual Conference, May 5-7, 2021

Conflict of interest Disclosure: I have no conflicts of interest



E-mail: hsiuju.chang@bccdc.ca





Existing disparities in sexual health service access experienced by sexual and gender minorities may have been exacerbated by public health responses to the COVID-19 pandemic

Objective: To describe differences in service access among a diverse sample of sexual health service clients in British Columbia (BC)



"Sex in the Time of COVID-19" Survey

- Anonymous, online survey
- Recruited participants from BCCDC's STI clinic & GetCheckedOnline clients ≥ 16 years old
- Conducted July 21 Aug 4, 2020

Survey Participant Sexual Health Needs by Gender and Sexual Orientation

Category: Gender and sexual orientation	# of participants	<pre># with sexual health need (%)</pre>
Men, straight	247	93 (38%)
Men, sexual minority	331	239 (72%)
Women, straight	342	196 (57%)
Women, sexual minority	221	139 (63%)
Non-binary, Genderfluid, Other gender	55	38 (69%)



Among participants who had a sexual health need:	Men, straight (N=93)	Men, sexual minority (N=239)	Women, straight (N=196)	Women, sexual minority (N=139)	Non-binary, Genderfluid, Other gender (N=38)
Accessed all needed services	35 (38%)	139/239 (58%)	101 (52%)	48 (35%)	17 (45%)
Barriers in accessing sexual health care during the COVID-19 pandemic					
There was public messaging that I was not supposed to seek healthcare that wasn't urgent	29 (31%)	64 (27%)	61 (31%)	67 (48%)	18 (47%)
I was concerned about getting COVID-19 while at a clinic or lab	22 (24%)	69 (29%)	53 (27%)	56 (40%)	22 (58%)
I was concerned about getting COVID-19 while travelling to a clinic or lab	19 (20%)	43 (18%)	34 (17%)	42 (30%)	18 (47%)
The place I usually go to for testing/care was closed/had reduced services because of COVID-19	22 (24%)	65 (27%)	50 (26%)	35 (25%)	10 (26%)
I didn't know where to access sexual health or STI testing services	12 (13%)	24 (10%)	23 (12%)	19 (14%)	3 (8%)
I was worried that a healthcare provider might judge me for having sex during COVID	8 (9%)	31 (13%)	21 (11%)	19 (14%)	6 (16%)
I live or am in close contact with someone at risk of COVID-19	3 (3%)	10 (4%)	5 (3%)	11 (8%)	11 (29%)

Bold: Three most prevalent barriers reported per gender/sexual orientation group



 The most commonly reported barriers to accessing sexual health services during the COVID-19 pandemic differed by gender and sexual orientation

 Inequities in sexual health care access experienced by sexual and gender minorities must be addressed in service planning and delivery throughout and beyond the pandemic response